



TRANSCELL ONCOLOGIS

Portfolio offering cell based platforms as preclinical research tools

TRANS-UC
(MESENCHYMAL STEM CELLS)



PRODUCT SHEET

1. **Intended Use:** This product is intended for preclinical application. It is not intended for any therapeutic or diagnostic use
2. **Storage Temperature:** -150°C or below
3. **Biosafety level to be used while formulating:** 1
4. **Description & Certificate of Analysis**

Product Information:

Catalogue Name	TRANSCHYMAL-UC
Catalogue Number	TRANS-UC
Source	Discarded Human Umbilical Cord Tissue
Special Handling	Minimize exposure to non-liquid nitrogen temperatures, especially room temperature transfers

Required Testing & Results:

Test	Specification	Result
Mycoplasma	Negative	Negative
Appearance, Container Integrity	Conforms	Conforms
Cell Count	> 1.0x10 ⁶ viable cells/ml	>1.2x10 ⁶ viable cells/ml
Cell Viability	>90%	>90%
Endotoxin	<3 EU/mL	<3 EU/mL

Bio Burden – Sterility	Negative	Negative
------------------------	----------	----------

Supporting Data:

Safety Data	Representative Final Product lots were found non-tumorigenic and non-toxic
Donor Specimen (DS)	This DS has met all release criteria including donor selection criteria, consent for umbilical cord blood donation, virus testing (HIV1/2, HBV & HBC)

General Information:

Organism	Homo sapiens, human
Source	Umbilical Cord Tissue
Cell type	Mesenchymal stem cells
Disease	Normal
Ethnicity	Indian
Product Format	Frozen

5. Safety Precaution

Recommending protective gloves, apron, face mask be worn when handling frozen vials

6. Unpacking & Storage Instructions

- Check all containers for leakage or breakage
- Remove the frozen vial from the dry ice packaging and immediately place at a temperature below -150°C, until ready for use

7. Handling Procedure for Frozen Cells

- Refer to the batch specific information for the total number of viable cells recovered from this lot
- Remove one vial at a time from storage and thaw the vial by gentle agitation in a 37°C water bath. Thawing should be rapid (1 to 2 minutes)

- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol under aseptic conditions
- Add the appropriate volume (5 mL) of medium into a sterile conical tube. Using a sterile pipette, transfer the cells from one cryo vial to the conical tube and gently tap the tube to mix the suspension for 5 min at room temperature
- Centrifuge the cells at 150 x g for 3 to 5 minutes
- Remove the supernatant and the pellet is to be re-suspended in the volume for desired next step

8. Human Material Precaution

All tissues used for isolation are obtained under informed consent and concerned regulatory ethics approval to conform to handling biological samples standards; to protect the privacy of the donor's personal health information

9. Transtoxbio Warranty

The viability of the product is warranted for 25 days from the date of shipment, and is valid only if the product is stored and handled according to the information included in

10. Disclaimer

- This product is intended for preclinical research purposes only and not for use in humans. While Transtoxbio uses realistic efforts to include support information on this product sheet, makes no warranties or representations as to its accuracy
- This product is sent with the condition that you are responsible for its safe storage, handling, and use. Transtoxbio is not liable for any damages or injuries arising from receipt and/or use of this product

11. Application Notes

TRAN-MSC is an *invitro* human sourced progenitor/stem cell based platform model composed of undifferentiated cells with self-renewing and differentiating capabilities. Recommended for 3D organ (bone & cartilage, wound) models. Ready to use; No culturing procedure involved to apply with customized configuration.